arXiv:1406.2550v1 [math.GR] 10 Jun 2014

A ONE-RELATOR GROUP WITH LONG LOWER CENTRAL SERIES

ROMAN MIKHAILOV

Abstract. A one-relator group with lower central series of length ω^2 is constructed. This answers a problem of G. Baumslag.

1. Introduction

For a group G, the lower central series are defined inductively as follows:

$$\gamma_1(G) = G,\ \gamma_{\alpha+1}(G) = [\gamma_\alpha(G), G]$$

and $\gamma_\tau(G) = \cap_{\alpha<\tau}\gamma_\alpha(G)$ for a limit ordinal τ. The smallest ordinal α, such that $\gamma_\alpha(G) = \gamma_{\alpha+1}(G)$ is called the *length of the lower central series of* G. First examples of finitely presented groups with the lower central series of length greater than ω were constructed by J. Levine [3]. T. Cochran and K. Orr constructed examples of 3-manifold groups with lower central series of length greater than ω in [2].

G. Baumslag asked the following question ([1], Problem 10): *Is the lower central series of a 1-relator group of length at most ωn for some finite ordinal n?*

Consider the following one-relator group

$$G = \langle a, b \mid a^{b^2} = aa^{3b}\rangle$$

This group is a semidirect product $F_2 \rtimes \mathbb{Z}$, where a generator of the cyclic group acts on $F_2 = F(x, y)$ as the automorphism

$$x \mapsto y$$
$$y \mapsto xy^3$$

The main result is the following

Theorem 1. *The group G has the lower central series of length ω^2.*

The group G provides an example of a one-relator group with length greater than ωn for any finite n, thus answering the above problem of G. Baumslag.

2. Proof of theorem 1

Let Γ be a group and M a $\mathbb{Z}[\Gamma]$-module. M is called *residually nilpotent* if

$$\cap_k M\Delta^k(\Gamma) = 0,$$

where $\Delta^k(\Gamma)$ is the kth power of the augmentation ideal $\Delta(\Gamma) := \ker\{\mathbb{Z}[\Gamma] \to \mathbb{Z}\}$.

This research is supported by the Chebyshev Laboratory (Department of Mathematics and Mechanics, St. Petersburg State University) under RF Government grant 11.G34.31.0026, and by JSC Gazprom Neft, as well as by the RF Presidential grant MD-381.2014.1.

Lemma 2. *Let M be a finitely generated free abelian group of rank $m \geq 1$ and an infinite cyclic group with generator t acts on M as a matrix $A \in GL_m(\mathbb{Z})$. Suppose that the product of any collection of eigenvalues of $A - Id$ is not ± 1. Then the module M is residually nilpotent $\mathbb{Z}[\langle t \rangle]$-module.*

Proof. Let Δ be the augmentation ideal of $\mathbb{Z}[\langle t \rangle]$. Observe that, for any $n \geq 1$, the power Δ^n is the ideal generated by $(1-t)^n$. Consider the submodule $N := M\Delta^\omega = \cap_{k \geq 1} M\Delta^k$ and assume that $N \neq 0$. The Krull intersection theorem implies that $NI = N$, hence $N(1-t) = N$ and the restriction of $(1-t)$ on N defines a bijection on N. Hence $det(A - Id) = \pm 1 = \beta_1 \dots \beta_k$, where $(\beta_1, \dots, \beta_k)$ is some collection of eigenvalues of $A - Id$. This gives a needed contradiction. Hence $N = 0$. □

Now we prove theorem 1. As usual, if $x, y, a_1, \dots, a_{k+1}$ are elements of a group G we set $[x, y] = x^{-1}y^{-1}xy$, $x^y = y^{-1}xy$ and define

$$[a_1, a_2, \dots, a_{k+1}] = [[a_1, \dots, a_k], a_{k+1}] \ (k > 1).$$

For $n \geq 1$, define T_n as a normal closure in G of all elements of the form $[x_1, \dots, x_n]$, where all x_i-s are either a or a^b. We have

$$H_1 = \langle a \rangle^G = F(a, a^b), \ H_2 = \langle [a, a^b] \rangle^G, \ H_3 = \langle [a, a^b, a], [a, a^b, a^b] \rangle^G$$

etc. Observe that

$$H_n = \gamma_n(H_1).$$

We claim that, for all $n \geq 1$,

$$\gamma_{n\omega}(G) \subseteq H_{n+1} \tag{1}$$

First we prove (1) for $n = 1$. Consider the quotient of G by the normal closure of the element $[a, a^b]$:

$$H = \langle a, b \mid a^{b^2} = aa^{3b}, [a, a^b] = 1 \rangle$$

The group H is the metabelian polycyclic group $(\mathbb{Z} \oplus \mathbb{Z}) \rtimes \mathbb{Z}$, where the generator of the cyclic group $\mathbb{Z}$ acts on $\mathbb{Z} \oplus \mathbb{Z}$ as the matrix

$$U := \begin{pmatrix} 0 & 1 \\ 1 & 3 \end{pmatrix} \tag{2}$$

The eigenvalues of the matrix $U - Id$ are $\frac{3 \pm \sqrt{13}}{2} - 1$. By lemma 2, the group H is residually nilpotent. Lets show now that $[a, a^b] \in \gamma_\omega(G)$. Observe that the element a is a generalized 3-torsion element, i.e. for every n there exists $t(n)$, such that $a^{3^{t(n)}} \in \gamma_n(G)$. Indeed, since

$$a^3 = [a, b^2]^{b^{-1}}, \tag{3}$$

$$a^{3^k} \in \gamma_{k+1}(G), \ k \geq 1.$$

Therefore, the element $[a, a^b]$ also is a generalized 3-torsion element. On the other hand, taking the commutator with a^b of the both sides of the relation, we get

$$[a^{b^2}, a^b] = [a, a^b]^{a^{3b}},$$

hence $[a^b, a] = [a, a^b]^{a^{3b}b^{-1}}$ and

$$[a^b, a]^2 = [a, a^b, a^{3b}b^{-1}] \tag{4}$$

Therefore $[a, a^b]$ is a generalized 2-torsion element. This implies that $[a, a^b] \in \gamma_\omega(G)$. Since H is residually nilpotent, we conclude that $\gamma_\omega(G) = \langle [a, a^b] \rangle^G$,

$G/\gamma_\omega(G) = H$ and the statement (1) is proved for $n = 1$. It is easy to see at this point that the group G has length greater than ω. Indeed, since $H_2(H) = \mathbb{Z}/2$ and $H_2(G) = 0$, we get

$$H_2(H) = H_2(G/\gamma_\omega(G)) \simeq \gamma_\omega(G)/\gamma_{\omega+1}(G) \simeq \mathbb{Z}/2$$

and

$$[a, a^b] \in \gamma_\omega(G) \setminus \gamma_{\omega+1}(G),\ [a, a^b]^2 \in \gamma_{\omega+1}(G).$$

Now suppose that the statement (1) holds for a given n. Here we prove this statement for $n + 1$.

We have an inclusion

$$\gamma_{n\omega+k}(G) \subseteq [H_{n+1}, \underbrace{G, \dots, G}_{k}],\ k \geq 1$$

Define the group $T := G/H_{n+2}$. By the assumption of induction, the term $\gamma_{n\omega}(T)$ lies in the quotient H_{n+1}/H_{n+2}, the term $\gamma_{n\omega+k}(T)$ is a subgroup of the quotient $([H_{n+1}, \underbrace{G, \dots, G}_{k}]H_{n+2})/H_{n+2}$ and can be presented as a product of conjugates of the elements

$$[\alpha, \underbrace{b, \dots, b}_{k}].H_{n+2},\ \alpha \in H_{n+1}.$$

Consider the quotient H_{n+1}/H_{n+2} as a $\mathbb{Z}[\langle b\rangle]$-module. If the module H_{n+1}/H_{n+2} is residually nilpotent $\mathbb{Z}[\langle b\rangle]$-module, i.e.

$$0 = \cap_k (H_{n+1}/H_{n+2})\Delta^k,$$

where Δ is the augmentation ideal of $\mathbb{Z}[\langle b\rangle]$, then the intersection

$$\cap_k ([H_{n+1}, \underbrace{G, \dots, G}_{k}]H_{n+2})/H_{n+2}$$

is trivial in T and the statement (1) will follow for $n + 1$. Denote $M := H_1/H_2$. Since H_1 is a free group and $H_i = \gamma_i(H_1),\ i \geq 1$, the quotient H_{n+1}/H_{n+2} is the $n+1$-st Lie power $L^{n+1}(M)$ by Magnus-Witt theorem (see [4], [6]). Lie powers are subgroups of tensor powers $L^{n+1}(M) \subseteq \otimes^{n+1}(M)$ and therefore,

$$\cap_k L^{n+1}(M)\Delta^k \subseteq \cap_k \otimes^{n+1}(M)\Delta^k$$

Here we consider the tensor powers of M as $\mathbb{Z}[\langle b\rangle]$-modules with the diagonal action of $\langle b\rangle$.

For given matrices A and B, the eigenvalues of their tensor product $A \otimes B$ consist of all possible products of the eigenvalues of A and B. The eigenvalues of the matrix U are $\alpha_1 = \frac{3+\sqrt{13}}{2}$, $\alpha_2 = \frac{3-\sqrt{13}}{2}$. Hence, the eigenvalues of the the tensor power $U^{\otimes n+1}$ are $\pm\alpha_1^l$ and $\pm\alpha_2^s$ for some $l, s \geq 1$. Suppose that one can present ± 1 as a product of elements of the form $\pm\alpha_1^l \pm 1$ and $\pm\alpha_2^s \pm 1$. Then the product of their conjugates in the sense of quadratic irrationality also is ± 1. Since $\alpha_1^l \pm 1$ is the conjugate quadratic irrationality to $\alpha_2^l \pm 1$, one can present ± 1 as a product of integers of the form

$$(\alpha_1^l - 1)(\alpha_2^l - 1) \text{ and } (\alpha_1^s + 1)(\alpha_2^s + 1)$$

for different $l, s \geq 1$. Since $(\alpha_1 - 1)(\alpha_2 - 1) = -3$ divides $(\alpha_1^l - 1)(\alpha_2^l - 1)$ for any $l \geq 1$, the absolute values of the terms $(\alpha_1^l - 1)(\alpha_2^l - 1)$ are greater than 1. For an odd s, the product $(\alpha_1^s + 1)(\alpha_2^s + 1)$ divides $(\alpha_1 + 1)(\alpha_2 + 1) = 3$. For an even s,

the terms (α_1^s+1) and (α_2^s+1) are greater than 1. We get a contradiction with assumption that one can present ± 1 as a product of elements of the form $\pm\alpha_1^l \pm 1$ and $\pm\alpha_2^s \pm 1$.

We conclude that, for any n, the matrix $U^{\otimes n+1} - Id$ satisfies the conditions of lemma 2 and therefore the $(n+1)$st tensor power $M^{\otimes n+1}$ is a residually nilpotent $\mathbb{Z}[\langle b\rangle]$-module. This implies that the Lie power $L^{n+1}(M)$ also is a residually nilpotent $\mathbb{Z}[\langle b\rangle]$-module and the inductive step is done,

$$\gamma_{(n+1)\omega}(G) \subseteq H_{n+2},$$

proving (1) for all n.

Since H_1 is a free group, it is residually nilpotent, and we get

$$\gamma_{\omega^2}(G) = \cap_n \gamma_{n\omega}(G) \subseteq \cap_n H_{n+1} = \cap_n \gamma_{n+1}(H_1) = 1.$$

Since $[a^b, a] \in \gamma_\omega(G)$, we have

$$1 \neq [[a^b,a],a,\underbrace{[a^b,a],\ldots,[a^b,a]}_{k}] \in \gamma_{(k+1)\omega}(G),\ k \geq 1.$$

This element clearly is non-trivial, since it is a basic commutator in the basic elements a, a^b of the free group $H_1 = F(a, a^b)$. Hence, the lower central series length of G is ω^2 and theorem 1 is proved.

Remark 3. *Observe that one can easily use the proof of theorem 1 to prove the following more general result. Let F_n be a free group and an element t acts on F_n as an automorphism $\phi : F_n \to F_n$. Define $\Phi = F_n \rtimes \langle t\rangle$. Suppose that any tensor power $(F_{ab})^{\otimes k}$ $(k \geq 1)$ is a residually nilpotent $\mathbb{Z}[\langle t\rangle]$-module. Then $\gamma_{\omega^2}(\Phi) = 1$. In particular, if Φ is not residually nilpotent, then the lower central series length of Φ is exactly ω^2.*

We add the next remark here due to its applications in the theory of localizations [5].

Remark 4. *The 2ωth lower central series term of G can be described as follows*

$$\gamma_{2\omega}(G) = \langle [a,a^b,a],[a,a^b,a^b]\rangle^G \tag{5}$$

We proved before that there is an inclusion $\gamma_{2\omega}(G) \subseteq H_3$. To see the converse inclusion, observe that a and a^b are generalized 3-torsion elements, hence

$$[a,a^b,a]^{3^l}, [a,a^b,a^b]^{3^l} \subseteq \gamma_{\omega+l}(G),\ l \geq 1.$$

However, the same is true for powers of 2, due to (4). Therefore, $H_3 \subseteq \gamma_{2\omega}(G)$ and description (5) follows.

Acknowledgements. The author thanks G. Baumslag, S.O. Ivanov, F. Petrov and K. Orr for discussions related to the subject of the paper.

References

[1] G. Baumslag: Some problems on one-relator groups; Proceedings of the Second International Conference on the Theory of Groups: Australian National University, August, 1973, *Lecture Notes in Mathematics*, **372** (1974).

[2] T. Cochran and K. Orr: Kent E. Stability of lower central series of compact 3-manifold groups, *Topology* **37** (1998), 497–526.

[3] J. Levine: Finitely-presented groups with long lower central series, *Israel J. Math.* **73**, (1991), 57–64.

[4] W. Magnus: Über Beziehungen zwischen höheren Kommutatoren, *J. reine angew. Math.* **177** (1937), 105-115.
[5] R. Mikhailov: On transfinite nilpotence of the Vogel-Levine localization, preprint
[6] E. Witt: Treu Darstellung Liescher Ringe, *J. reine angew. Math.* **177** (1937), 152-160

Chebyshev Laboratory, St. Petersburg State University, 14th Line, 29b, Saint Petersburg, 199178 Russia and St. Petersburg Department of Steklov Mathematical Institute

E-mail address: rmikhailov@mail.ru